
Taking Inventory of the Universe: The Mystery of Dark Matter

Stanley Seibert
University of Pennsylvania
LBNL, February 20, 2013




Today's topic: A gravitational mystery...

Abell 1703

...brought to you by precision astronomy

Seven Decades of "Excess Gravitation"









Rotation Curves

Gravitational Lensing


Cluster Collisions



CMB Power Spectrum



Baryon Acoustic Oscillations



Simulations of Structure Formation

And many others!


Responses to the Unexpected


- I."Observational" error.
 - "Precision astronomy is hard. Maybe you made a mistake."
- 2. Interpretation or modeling error.
 - "Are you really sure you know where all the baryons are?" (Black holes, neutron stars, brown dwarfs, etc.)
- 3. New or modified interaction between matter. *Modified Newtonian Dynamics (MOND)*, TeVeS
- 4. New material constituent. (usually a new particle) Neutralinos, axions, Kaluza-Klein states, ...

Pick any combination of all 4!

The Dark Matter Hypothesis


A substantial fraction of the matter in the universe is in a form that rarely (or never) interacts with photons, rendering it invisible ("dark") to direct electromagnetic observation.






But, isn't proposing a new form of matter based purely on astronomical evidence preposterous?


We've Been Here Before...





E. Frankland N. Lockyer

Helium was first discovered by astronomers in the solar chromosphere in 1868, but not by chemists in the lab until 1895!



W. Ramsay

What is Dark Matter?

Suppose you decide to search for "terrestrial" dark matter. What do you know?

If you explain the astronomy data with dark matter, then you know:

- Cross-sections for interaction between dark matter and itself/other particles are very small.
 (or you would have seen it already)
- Local density near Earth is around 0.3 GeV/cm³ (within a factor of 2 or 3)
- Solar system moves through dark matter halo at 220 km/sec



Direct Dark Matter Searches


("looking for your lost keys under the street light")

- Anomalous nuclear recoils (WIMP scattering)
- XENON, CDMS, CoGeNT, DEAP/ CLEAN, LUX, PICASSO, COUPP, CRESST, XMASS, EDELWEISS, ...
- 2. Primakoff interactions (axion-photon coupling)

ADMX, CAST, ...

- 3. Periodicity/Directionality DAMA/LIBRA, DRIFT, DMTPC, ... (the 21st century search for the "aether wind")
- 4. [Insert your clever idea here]


Why WIMPs?



A new particle with mass near the EW symmetry-breaking scale and weak force gauge couplings produces the right thermal relic density.

Annu. Rev. Astron. Astrophys. 2010. 48:495-545

Hunting for WIMPs




Want:

Low energy threshold

Large, "cheap," and clean target material


Excellent separation of nuclear recoils (induced by WIMPs) from other backgrounds

Background Discrimination



Experimental Results: How are we doing so far?

In the Thick of It




Summary of the data so far:

It took astronomers many decades to sort out evidence for dark matter.

Particle physicists are still very early in the process, and there is plenty of room for other approaches....

Can we look to past successes for inspiration?

Lessons From Neutrino Experiments



Low energy neutrino detection demands large, clean, and deep underground experiments, much like dark matter.


Scaling requires a detector that can be composed from "simple" repeatable structures.

Percent-level calibration is possible, especially when you start with a well-modeled detector.

Careful modeling of backgrounds allows for low threshold signal extraction.


Sudbury Neutrino Observatory

Ex: SNO Radon Spike




Best fit convolution of Monte Carlo by ~0.1 MeV

MiniCLEAN: Searching for Dark Matter with Argon and Neon Scintillation




Scintillation in Noble Liquids




Energy deposition in noble liquids produces short lived excited diatomic molecules in singlet and triplet states.

Pulse Shape Analysis



Rejecting Electron-like Events in Argon




Discriminate with ratio of prompt to total light

Reject beta and gamma backgrounds with less than 10-8 leakage


Important to reject intrinsic ³⁹Ar background

Observing Extreme UV



Almost everything absorbs 128 nm light! TPB can wavelength shift EUV up to 440 nm with high efficiency.


TPB Re-emission Efficiency



10% measurement resolves factor of 3 ambiguity in the literature

→ Disentangles TPB and argon scintillation efficiency, useful for other noble liquids

TPB Re-emission Spectrum



NIM A 654 (2011), 116-121 V. Gehman, SRS, K. Rielage, A. Hime, Y. Sun, D.-M. Mei

Discovered that re-emission spectrum independent of illumination wavelength and has cutoff near 400 nm.

→ Direct impact on choice of optical materials

Single Phase Ar/Ne Detectors

Advantages:

- Target material is very inexpensive.
- No need for electric fields to drift charge.
 - Simpler detector design
 - Able to use a spherical geometry (but not required)
 - Does not require ³⁹Ar-depleted argon for large detectors
- Neon is clean enough to use for pp solar neutrinos

Disadvantages:

- Lower A² reduces coherent scattering enhancement
- Self-shielding from external backgrounds worse than other materials
- Atmospheric argon contains a high rate beta decay isotope, ³⁹Ar @ I Bq/kg (also a fantastic calibration!)

The DEAP and CLEAN Family of Detectors

10⁻⁴⁴ cm²

 10^{-45} cm^2

 10^{-46} cm^2

WIMP or Sensitivity

DEAP-0:

Initial R&D detector

DEAP-I:

7 kg LAr2 warm PMTsAt SNOLab since 2008

DEAP-3600:

3600 kg LAr (1000 kg fiducial mass) 266 warm PMTs SNOLAB 2014

picoCLEAN:

Initial R&D detector

microCLEAN:

4 kg LAr or LNe2 cold PMTssurface tests at Yale

MiniCLEAN:

500 kg LAr or LNe (150 kg fiducial mass) 92 cold PMTs SNOLAB 2013

40-140 tonne LNe/LAr Detector:

pp-solar V, supernova V, dark matter < 10⁻⁴⁶ cm² ~2018?



MiniCLEAN Collaboration



Boston University

D. Gastler, E. Kearns, S. Linden

UC Berkeley

G.D. Orebi Gann

Los Alamos National Laboratory

M. Akashi-Ronquest, K. Bingham, R. Bourque, J. Griego, A. Hime, F. Lopez, J. Oertel, K. Rielage, L. Rodriguez

Massachusetts Institute of Technology

J.A. Formaggio, S. Jaditz, J. Kelsey, K. Palladino

National Institute Standards and Technology

K. Coakley

University of New Mexico

M. Bodmer, F. Giuliani, M. Gold, D. Loomba, J. Matthews, P. Palni, J. Wang

University of North Carolina/TUNL

R. Henning

University of Pennsylvania

T. Caldwell, J.R. Klein, A. Mastbaum, S. Seibert

Royal Holloway University of London

A. Butcher, J. Monroe, J.A. Nikkel, J. Walding

University of South Dakota

V. Guiseppe, D.-M. Mei, G. Perumpilly, C. Zhang


Syracuse University

R.W. Schnee, B. Wang

Yale University

D.N. McKinsey

The MiniCLEAN Detector



92 8" PMTs


TPB @ R=43 cm

PMTs @ R=81 cm


Modular "Cassettes"




Simplified View



Water Shielding




SNOLAB



Lab Area




Inner Vessel




Manufacturing completed, pressure and leak tested in Sept. 2012

Inner Vessel Underground




MiniCLEAN WIMP Analysis

Will perform a maximum likelihood analysis with a blind signal box in three reconstructed observables:



WIMP Sensitivity



MiniCLEAN w/ 150 kg fiducial volume

80 = 80 keVr threshold

50 = 50 keVr threshold

30 = 30 keVr threshold

Small change in energy threshold equivalent to large change in fiducial volume!

Lowering the Energy Threshold


- Rejection of ³⁹Ar beta decay events sets the energy threshold in single-phase argon.
- ³⁹Ar is the only background that scales like volume, rather than surface area, so it also limits detector size.
- Important to maximize the performance of our pulse-shape rejection algorithms.
- Prior to construction, we have done this with a detailed simulation...

MiniCLEAN Simulation: RAT




- Developed for Braidwood (SRS), now used by SNO+ and DEAP/CLEAN.
- Wraps together GEANT4, ROOT, and GLG4sim (KamLAND) into full simulation and analysis package.
- Fully propagates optical photons, including PMT response and digitizer / DAQ simulation
- Includes pulse shape analysis tools, maximum likelihood position reconstruction

Comparison with MicroCLEAN



Modeling of Cryogenic PMT Response




MiniCLEAN uses Hamamatsu R5912-02-MOD 8" PMTs, tested for use at temperatures down to 25K.

Photoelectron (PE) Counting

- Counting the number of PE on a channel is fundamental to energy and position estimation.
- PMTs produce variable size, finite width (~20ns) pulses for each PE.
- Counting peaks in a waveform is biased low when there is pulse pileup.
- Integrating charge is unbiased, but higher variance.
- How to get best of both worlds?
- (Side note: This is one of many instances where single phase detectors get better as they get bigger and add channels...)

Bayesian PE Counting



Bayesian PE Counting

Chop up waveform for each PMT into regions with pulses, and for each region, compute:

 $P_N(n|q,t_1,t_2) = \underbrace{\frac{P_Q(q\,|\,n)P_N(n\,|\,t_1,t_2)}{P_Q(q\,|\,i)P_N(i\,|\,t_1,t_2)}}_{n\text{-PE charge distribution}}$

Probability of *n*-PE given pulse integral *q*, and pulse ranging from t₁ to t₂

Poisson probability of observing n (or i) photons between t_1 and t_2 in this PMT.



This is where we insert our knowledge of argon scintillation time structure and event position & energy

Position & Energy Reconstruction



No photon can travel directly from the event vertex to a PMT!


- Rayleigh scattering length in argon is ~90 cm (or 66 cm?).
- TPB further randomizes photon paths.
- Estimating PMT hit probability based simply on solid angle tends to bias the fit inward.
- The actual light pattern is more isotropic than a direct propagation model would predict.

ShellFit: Self-Tuning Maximum Likelihood Fitting

- Automatically use Monte Carlo simulation to collapse complete space of photon histories to before and after wavelength shifting.
- Integrate over TPB surface using a GPU to compute likelihood.

 4 sec/event w/ CPU and 0.2 sec/event w/ GPU! UV transmission to TPB Visible


Energy resolution after iterating PE counting




MiniCLEAN simulation, ³⁹Ar, 75-100 PE, inside fiducial volume

Particle ID Techniques

Prompt ratio (standard)




Likelihood ratio (new)




The Future

 The "Generation 2" concept for single phase liquid argon and neon is the CLEAN experiment.



The Future

Simpler PMT modules



(Front TPB plate removed)

The Future

- Between 40 and 140 tons total argon mass (≥15 tons fiducial).
- Optimal size with natural argon limited by pulse-shape discrimination performance and pileup.
- At ~140 tons (total), a dual-use argon and neon detector can do a percent-level precision measurement of pp solar neutrinos and observe supernova neutrinos.
- With depleted argon, could go very large...

Conclusion

- Something is out there, and it might be WIMPs!
- We've seen claims of direct detection, but you should continue to be skeptical.
- Single phase noble liquid detectors offer a highly scalable option for dark matter and neutrino detection.
- MiniCLEAN extends the DEAP/CLEAN series of detectors to 150 kg fiducial volume with liquid argon and neon.
- Many analysis improvements ready to test on data as it arrives.
- Construction is underway, with detector commissioning scheduled for mid-2013.

"I often look at the bright yellow ray emitted from the chromosphere of the sun, by that unknown element, Helium, as the astronomers have ventured to call it. It seems trembling with excitement to tell its story, and how many unseen companions it has. And if this be the case with the sun, what shall we say of the magnificent hosts of the stars? May not every one of them have special elements of their own? Is not each a chemical laboratory in itself?"

John William Draper Inaugural Address to the American Chemical Society 1876