


The Universe as a fundamental Physics laboratory




Neutrinos

Masses? Hierarchy? Dirac/Majorana? Additional sterile neutrinos? CP phase?


Cosmology can weigh the neutrinos

Neutrinos = 0.5% of all matter, but their gravity suppresses LSS 8-fold = 4%



→ CMB & LSS: masses to ~ 20 meV precision

Why now? LSS experiments are probing dark matter, dark energy, neutrinos & the Universe's initial conditions






Dark Energy Spectroscopic Instrument

Vera Rubin Observatory LSST

SPHEREX

Why care about baryons? Cosmology & Galaxy formation




Illustris


Supernovae and supermassive black holes regulate galaxy formation Unknown feedback amplitude

→ Missing "baryon problem"


How to analyze 1% precision LSS data when baryons (15% of matter) are missing?




Starlight




Starlight, dark matter



Starlight, dark matter, gas density




Starlight, dark matter, gas density, temperature



Starlight, dark matter, gas density, temperature, velocity



Backlighting the galaxies with the CMB



Summary of CMB secondaries

Key parameters:

$$\theta_{\rm lens} \sim 1' \sim 10^{-4}, \ \int dt \dot{\Phi} \sim 10^{-4}, \ \tau \sim 10^{-3}, \ \frac{k_B T_e}{m_e c^2} \sim \left(\frac{v_{\rm th}}{c}\right)^2 \sim 0.01, \ \frac{v_{\rm bulk,rot,turb}}{c} \sim 10^{-3}, \ \frac{\delta T_0}{T_0} \sim a_2 \sim 10^{-5}$$

→ Many imprints with complementary information :

Potential

Lensing ISW, Rees-Sciama Moving lens

$$\begin{array}{cc} \theta_{\rm lens} \frac{\nabla T_0}{T_0} & {\rm Total\ mass} \\ \int dt\ \dot{\Phi} & {\rm DE,\ accretion\ rate} \\ \theta_{\rm lens} (v_{\rm bulk}\ \bot/c) & {\rm Transverse\ velocities} \end{array}$$

Single scattering

Screening kSZ, rot kSZ, turb kSZ tSZ, relat tSZ Polarized scattering


$$au~(\delta T_0/T_0)$$
 Gas density $au~(v_{
m bulk}~||/c)$ Gas density, LOS velocities $au~(v_{
m th}/c)^{2,4}$ Gas thermal pressure, temperature $au~(v_{
m bulk}~\perp/c)^2$, $au~a_2$ Gas density, Ultra large scales


Multiple scattering

Smaller by factor au

Break degeneracies with tau?

Why now? High-res high-sensitivity CMB experiments





SO Science goals and forecasts 19 Lee+20


Growing interest in the CMB





Outline: Combining CMB & LSS

Gas shadows






ACT x DESI kSZ & tSZ


Hadzhiyska+23, 24

RiedGuachalla+23, 25

Liu+24

Mass shadows





CMB lensing noise bias avoidance
Shen Schaan Ferraro 23

Bayesian lensing biasfrom polarized extragalactic foregrounds

Qu Millea Schaan 24


Gas shadows: Localize the missing baryons

Missing baryon "opportunity": Cosmology & Galaxy formation

Galaxy formation

Feedback pushes gas outside virial radius Too faint to detect

- → Missing baryon problem
- → Limits our understanding of galaxy formation




Haider+16, Illustris simulation


Cosmology

Baryons ~15% total matter

- → Largest (30%) theoretical uncertainty on matter power spectrum
- → Limits cosmology from weak lensing: Rubin, Roman, Euclid




CMB can help: Sunyaev-Zel'dovich effects




→ Unique probe of missing baryons! (Looking forward to FRBs too!)

Extracting tSZ: single galaxy



SNR per galaxy is too low to detect!

Extracting tSZ: 400,000 galaxies




Extended tSZ profile is well resolved!


Higher precision tSZ: ACTxDESI

Henry Simone Emmanuel
Liu Ferraro Schaan

Total Control of the con

High precision stacked tSZ
Significant dust bias
Clean by masking/deprojecting with variable dust SED






Schaan Ferraro+20

Higher precision tSZ: ACTxDESI

Henry Simone Emmanuel
Liu Ferraro Schaan


Total Control of the Con

High precision stacked tSZ
Significant dust bias
Clean by masking/deprojecting with variable dust SED



Liu+24


CMB can help: Sunyaev-Zel'dovich effects



→ Unique probe of missing baryons! (Looking forward to FRBs too!)

Extracting kSZ: Velocity reconstruction

Mass conservation: $\vec{v} \propto \vec{\nabla} \Delta^{-1} \delta$



kSZ amplitude \sim < v_{true} v_{rec} > or $r(v_{true}, v_{rec})$ \rightarrow Quantify from sims


Velocity reconstruction via continuity equation: assessment


Bernardita Boryana Simone Emmanuel Ried Guachalla Hadzhiyska Ferraro Schaan











Hadzhiyska+23, Ried Guachalla+23


Key results

- Reconstruct halo (not galaxy) velocities
 - → good for kSZ!
- Photo samples extremely powerful (DESI LS, DES, HSC, LSST)
- Naïve hybrid photo-spectro reconstruction worse than photo-only

Assess impact of

- satellite fraction
- number density
- smoothing
- cosmology

Higher precision measurements: ACTxDESI



BOSS (CMASS AND LOWZ)

Higher precision kSZ: ACTxDESI


Bernardita Emmanuel Boryana Simone Ried Guachalla Schaan Hadzhiyska Ferraro









High precision stacked kSZ
Compare photo & spectro samples
Explore dependence on M, z, L
Confirm gas more extended than DM
Matches Illustris (large feedback),
not Illustris TNG





Ried Guachalla+25

Aren't CMB experiments too low resolution?



Ried Guachalla+25

Virial radius of galaxy groups $\sim 1'$, and gas extends several $R_{vir} \rightarrow Well$ matched to CMB data!

kSZ unique probe of outskirts of galaxy groups and clusters


→ Complementary with X-ray & tSZ, which probe cluster centers best

Latest stacked kSZ profiles available?


DESI LRG, BGS, ELG

spectro: Ried Guachalla+25


photo: Hadzhiyska+25




Also splits by redshift,



stellar mass,




absolute magnitude



→ All publicly available. Don't hesitate to use them & ask questions!


Large-scale noise: profile slope VS Mgas




Primary CMB (~degree) dominates noise on larger scales


 \rightarrow kSZ measures profile shape/slope better than its integral (M_{gas})

CAP filter avoids displaying constant CMB mode, but is not required. Fourier measurements simplify this





kSZ is faint. Is precision enough for lensing?




Precision already informative in 2021, now much better & improving

Baryons are 15% of the matter.

For 1%-precision lensing, only 7%-precision kSZ needed

- → statistics will not be limiting
- → Modeling is the challenge

Subtracting baryons from galaxy-galaxy lensing is a well-posed problem



→ Directly subtract the baryonic contribution!

Same halos, HOD, weighting (linear in mass, VS tSZ or Xray), angular scales

Hadzhiyska+25, Sunseri+25, McCarthy+25

Galaxy formation & Cosmic shear: trickier!

kSZ only measured around some halos at some z Cosmic shear = sum over halo masses & z

→ Extrapolation needed (Lucie-Smith+25)

Universality of matter power suppression? (Van Daalen+20, Joop's talk)
→ would help!

Comparison with simulation requires matching galaxy sample


mass (stellar or halo? mean or distribution?) satellite fraction & miscentering HOD?

→ All "large feedback" claims hinge on this

Great progress being made in modeling Bigwood+24, Sunseri+25, McCarthy+25

Outline: Combining CMB & LSS

Gas shadows






ACT x DESI kSZ & tSZ


Hadzhiyska+23, 24

RiedGuachalla+23, 25

Liu+24

Mass shadows







CMB lensing noise bias avoidance
Shen Schaan Ferraro 23

Bayesian lensing biasfrom polarized extragalactic foregrounds


Qu Millea Schaan 24






~2' deflections, coherent on degree scale


Surface brightness: $T(\mathbf{x}) = \tilde{T}^0(\mathbf{x} - \mathbf{d}_{(\mathbf{x})})$ Born approximation: $\mathbf{d} = \nabla \phi = 2\nabla \Delta^{-1} \kappa$




~2' deflections, coherent on degree scale Surface brightness: $T(\mathbf{x}) = T^0(\mathbf{x} - \mathbf{d}_{(\mathbf{x})})$ Born approximation: $\mathbf{d} = \nabla \phi = 2\nabla \Delta^{-1} \kappa$



~2' deflections, coherent on degree scale Surface brightness: $T(\mathbf{x}) = T^0(\mathbf{x} - \mathbf{d}_{(\mathbf{x})})$ Born approximation: $\mathbf{d} = \nabla \phi = 2\nabla \Delta^{-1} \kappa$




~2' deflections, coherent on degree scale Surface brightness: $T(\mathbf{x}) = T^0(\mathbf{x} - \mathbf{d}_{(\mathbf{x})})$ Born approximation: $\mathbf{d} = \nabla \phi = 2\nabla \Delta^{-1} \kappa$



Lensing breaks the statistical isotropy of the CMB by coupling small and large scales

→ Reconstruct with a quadratic estimator

CMB lensing noise bias avoidance




Shen Schaan Ferraro 2024

Slides adapted from Delon Shen




Gaussian noise bias subtraction is limiting on small scales






Gaussian noise bias subtraction is limiting on small scales





Gaussian noise bias subtraction is limiting on small scales



Noise bias present even without lensing → Subtract it!




Recall
$$\hat{\kappa}_{\boldsymbol{L}} \sim \int_{\boldsymbol{\ell}} T_{\boldsymbol{\ell}} T_{\boldsymbol{L}-\boldsymbol{\ell}} \sim \sum_{\boldsymbol{\ell}} \sum_{\boldsymbol{\ell}} \sum_{\boldsymbol{k}} \sum_{\boldsymbol{k}}$$




Recall
$$\hat{\kappa}_{m{L}} \sim \int_{m{\ell}} T_{m{\ell}} T_{m{L}-m{\ell}} \sim \sum_{m{\ell}} \sum_{m{\ell} \in \mathbb{Z}_{+}} \sum_{m{\ell} \in \mathbb{Z}_{+}}$$

$$\langle \hat{\kappa}_{m{L}} \hat{\kappa}_{m{L}}^* \rangle \sim \sum_{ ext{Quadrilaterals}} L_{---}$$















Idea: ignore all the quadrilaterals that **do** contribute to the Gaussian bias, the parallelograms




Useful with mask & noise anisotropy






Useful with mask & noise anisotropy



Useful even for realistic inhomogeneous noise No computational cost No statistical cost


Bayesian lensing bias is robust to polarized extragalactic foregrounds &


Bayesian analog of point source hardening



Qu Milllea Schaan 2024

Extragalactic foregrounds: Temperature





Power dominant on small scales

→ extra noise

Statistics is non-Gaussian, uncertain and correlated with κ_{CMB}


- → Several percent bias (bispec.+trispec.)

 Schaan Ferraro 18
- → limits $\ell_{\text{max}} \approx 3000$ for SO and S4

Point source/profile hardening, shear, asymmetric QE help

Sailer+21, 23, Darwish+23, Madhavacheril Hill 18 & many more


Extragalactic foregrounds: Polarization




Fewer foregrounds, smaller power, simpler statistics (Poisson) Radio power scales as sensitivity after masking point sources Optimal methods outperform QE in polarization


→ Sensitivity to foregrounds? Analog of point source hardening?

Optimal lensing fits point sources as convergence dipoles







Optimal lensing fits point sources as convergence dipoles




Optimal lensing fits point sources as convergence dipoles




QE too!





Bias to Bayesian lensing is small for CMB-S4

Ratio of bias to statistical uncertainty




Qu Millea Schaan 24

For futuristic experiments where the bias may be important, we derive an analog to point source hardening

Outline: Combining CMB & LSS

Gas shadows






ACT x DESI kSZ & tSZ


Hadzhiyska+23, 24

RiedGuachalla+23, 25

Liu+24

Mass shadows





CMB lensing noise bias avoidance
Shen Schaan Ferraro 23

Bayesian lensing biasfrom polarized extragalactic foregrounds

Qu Millea Schaan 24

Conclusions



Key Astrophysics & BSM physics


Galaxy formation, dark energy, dark matter, inflation, neutrinos








Key new experiments coming along DESI, Rubin LSST, Simons Observatory





CMB is a large-scale structure probe

Useful insights from combining CMB & LSS

Thank you!